arXiv:1302.0478v1 [cond-mat.stat-mech] 3 Feb 2013

Fourier's Law and temperature gradient in a 1D harmonic lattice

K.V. Reich*

(Dated: February 5, 2013)

Heat flow and thermal profile in a 1D harmonic lattice with coordinate-dependent masses has been calculated in the thermodynamic limit. It is shown in the particular example of a 1D harmonic lattice with linearly increasing masses that in standard Langevin conditions of contact, a temperature gradient can form, and Fourier's law can be obeyed.

It is well known [1] that in one-dimensional harmonic systems no temperature gradient can evolve, and neither can Fourier's law be obeyed, i.e., the dependence of heat flux J on the size of a system N cannot be described by the relation $J \sim 1/N$.

This conclusion was drawn on the basis of a consideration of only two systems for which these results can be obtained with analytical accuracy, more specifically, systems of particles with equal or alternating masses [2, 3]. It came therefore as a surprise when numerical simulations demonstrated that in systems with linearly [4] or exponentially [5] increasing masses a temperature gradient can form.

Interest in 1D systems in which mass depends on coordinates did not come as a surprise. It stemmed primarily from studies of functional-gradient materials (FGM) which can be employed in devices designed for monitoring and control of heat flow on a nanoscale [5–8].

The present study demonstrates a method of solution of linear equations describing such functional-gradient materials. In particular, for a system with an arbitrary dependence of masses on coordinate an exact analytical solution has been derived, which describes the temperature profile and the heat flow. It will be shown that in a system with a linear mass distribution heat gradient does indeed exist, and Fourier's law is indeed obeyed in certain cases.

Consider a one-dimensional chain of $N + 1$ particles which interact harmonically with their nearest neighbors with a spring constant K. The momentum and displacement n of a particle of mass m_n will be denoted by p_n and x_n, respectively. The Hamiltonian of this system (Fig. 1) can be written in the form:

$$H = \sum_{n=0}^{N} \left(\frac{p_n^2}{2m_n} + \frac{1}{2} K (x_{n+1} - x_n)^2 \right), \tag{1}$$

The equations of motion for such a system become:

$$x_n \cos\beta_n = \frac{1}{2} \left(x_{n+1} + x_{n-1} \right), \tag{2}$$

where $\cos\beta_n = (1 - \omega^2 m_n / 2K)$.

In the thermodynamic limit the solution of this system of equations can be obtained in analytical form.

To do this, we use Fourier's method by writing x_n in the form



FIG. 1. Schematic of a harmonic system with coordinate-dependent masses

$$x_n = \int_{k_1}^{k_2} f(k) \exp(ikn) dk. \tag{3}$$

We choose the limits of integration k_1, k_2 such that the function $f(k)$ and all its derivatives tend to zero at these points. In this case we come to the equality

$$(-in)^p x_n = \int_{k_1}^{k_2} \frac{d^p f(k)}{dk^p} \exp(ikn) dk.$$

We further assume that the mass m_n and β_n are functions of n/N. Now the function $\cos\beta(n/N)$ can be expanded in the Taylor series with respect to the variable n/N, with Eqs. (2) converting into a differential equation for function f:

$$\cos\beta \left(i \frac{1}{N} \frac{d}{dk} \right) f = f \cos(k), \tag{4}$$

Only in rare cases, such as a linear dependence of particle masses on index:

$$m_n = M_0 + \frac{M_1 - M_0}{N} n, \tag{5}$$

this differential equation allows an exact solution.

Because to this case we are going to revert more than once, we note that Eqs. (2) are solved in terms of Hankel functions: $x_n = H^{(1)}_{n+\gamma}(z), H^{(2)}_{n+\gamma}(z)$, where $\gamma = z \cos\beta(0), z = N/(\cos\beta(1) - \cos\beta(0))$. This can be verified by direct substitution using the recurrent relation $2n/z Z_n(z) = Z_{n+1}(z) + Z_{n-1}(z)$, $Z_n(z) = H^{(1)}_n(z), H^{(2)}_n(z)$ [9].

In a general case, we are interested in the solution in the thermodynamic limit. We are going to look for the solution to Eq. (4) subject to the condition $N \to \infty$ in the form:

$$f(k) = \varphi(k) e^{ig(k)N}.$$

In this case we come to the following relations for g and φ:

$$g = -\int^k \beta^{-1}(z) dz, \tag{6}$$

$$\varphi(\beta(k)) = \exp\left(-\int^k \frac{dx}{M'(x)} \left(\frac{M(x) - M(0)}{x} \right)' \right), \tag{7}$$

where β^{-1} is a function inverse of β. Note that φ does not depend on frequency.

Next we choose the path of integration in Eq. (3) such that it will descend most steeply from the saddle point. Substituting the expressions for g and φ, we come in the limit that $N \to \infty$ to a particular solution to Eqs. (2):

$$y_n = \frac{1}{\sqrt{\sin\beta(a)}} \phi(a) e^{ib\beta(a)} e^{iN \int\limits_0^a \beta(x)dx}, \tag{8}$$

where n is taken in the form $n = aN + b$, $\phi(k) = \varphi(\beta(k))\sqrt{|M'(k)|}$.

We finally come to the general solution of Eqs. (2) in the form $x_n = Ay_n + By_n^*$.

Using the standard non-equilibrium Green function method [1, 10, 11] and assuming the temperature on the left to be $T_L = T + 1/2\Delta T$, and on the right, $T_R = T - 1/2\Delta T$, with $\Delta T, \omega_n \ll T$, we can obtain both the heat flux J in the system, and the temperature profile T_n:

$$\begin{aligned} J &= \frac{\gamma}{m_N} \Delta T I_N, \\ T_n &= T + \frac{1}{2} \Delta T (I_n - \bar{I}_n), \end{aligned} \tag{9}$$

where the quantities

$$\begin{aligned} I_n &= \frac{m_n \gamma}{\pi} \int\limits_{-\infty}^{\infty} d\omega \omega^2 |G_{0n}|^2, \\ \bar{I}_n &= \frac{m_n \gamma}{\pi} \int\limits_{-\infty}^{\infty} d\omega \omega^2 |G_{nN}|^2 \end{aligned} \tag{10}$$

are expressed in terms of the Green's function:

$$G(\omega) = \left(K - M\omega^2 - \Sigma_L(\omega) - \Sigma_R(\omega)\right)^{-1}.$$

with M being a diagonal matrix with elements corresponding to the particle masses, and K, the dynamic matrix for the system. The function $\Sigma(\omega)$ specifies the conditions of contact of the system under consideration with heat reservoirs. For the standard (Langevin) contact, $\Sigma_L(\omega) = i\omega\gamma\delta_{i,0}\delta_{j,0}$, $\Sigma_R(\omega) = i\omega\gamma\delta_{i,N}\delta_{j,N}$. After a few straightforward transformations similar to those made in Ref. [12], we come to:

$$G_{0,k} = \frac{\left|D_{k+1,N} - \Sigma D_{k+1,N-1}\right|^2}{\left|D_{0,N} - \Sigma(\omega)(D_{1,N} + D_{0,N-1}) + \Sigma(\omega)^2 D_{1,N-1}\right|^2}, \tag{11}$$

where $D_{l,m}$ is defined to be the determinant of the submatrix of $K - \omega^2 M$ beginning with the l-th row and column and ending with the m-th row and column. This determinant can be readily derived, because we know the general solution for Eqs. (2):

$$D_{lm} = \frac{\operatorname{Im} y_{l-1} y_{m+1}^*}{\operatorname{Im} y_{l-1} y_l^*} \tag{12}$$

The solution of equation $D_{0N} = 0$ with respect to frequencies ω yields the dependence of the wave vector k on the frequency of vibrations in the system under consideration $k(\omega)$:

$$Re \int\limits_0^1 \beta(x) dx = k(\omega) \tag{13}$$

In particular, for the case of linear mass distribution (5) we come to:

$$k(\omega) = \operatorname{Re} \frac{f(\beta(0)) - f(\beta(1))}{\Delta_{\cos}},$$

where $f(\beta) = (\sin\beta - \beta\cos\beta)$. This dispersion relation is displayed in Fig. 2. This result correlates with numerical simulations [13].

As seen from Fig. 2, in systems with linearly increasing masses one can identify two kinds of vibrations, more specifically, delocalized phonons with frequencies $\omega < \sqrt{2}\omega_0$, and localized "gradons" with frequencies $\sqrt{2}\omega_0 < \omega < \sqrt{2}\omega_1$, $\omega_{0,1} = \sqrt{2K/M_{0,1}}$.

In a general case, there exist also phonons with frequencies $\omega < \sqrt{2}\omega_{min}$ and gradons with frequencies $\omega > \sqrt{2}\omega_{min}$, with ω_{min} being determined by the maximum mass in the system. Incidentally, for gradons β becomes imaginary, with the determinant starting to grow exponentially. As a result, frequencies $\omega > \sqrt{2}\omega_{min}$ do not contribute to the integrals in Eqs. (10).

Standard methods of averaging [2] permit reducing the integrands in Eq. (10) in the $N \to \infty$ limit to the form

$$\begin{aligned} |G_{0N}|^2 &= \left(\frac{\phi(0)}{\phi(1)}\right)^2 \frac{\sin\beta(0)\sin\beta(1)}{\gamma\omega(1+\gamma^2\omega^2)(\sin\beta(0) + \sin\beta(1))} \\ |G_{0n}|^2 &= \frac{1}{2\gamma\omega} \left(\frac{\phi(0)}{\phi(x)}\right)^2 \frac{\sin\beta(0)}{(\sin\beta(0) + \sin\beta(1))\sin\beta(a)} \\ |G_{nN}|^2 &= \frac{1}{2\gamma\omega} \left(\frac{\phi(x)}{\phi(1)}\right)^2 \frac{\sin\beta(1)}{(\sin\beta(0) + \sin\beta(1))\sin\beta(a)} \end{aligned} \tag{14}$$



FIG. 2. Dispersion relation of the vibration energy in a harmonic chain with linearly increasing masses (5) plotted vs. the wave vector for different relations between the boundary masses $\omega_{0,1} = \sqrt{2K/M_{0,1}}$. For energies $\omega < \sqrt{2}\omega_0$, the system operates with delocalized phonons, and for $\sqrt{2}\omega_0 < \omega < \sqrt{2}\omega_1$, with localized gradons.

where n is used in the form $n = aN$. To preclude misunderstanding, we note that these relations do not go one into another at $a = 1$ and $a = 0$, because $N \to \infty$ is a non-uniform limit.

Putting relations (14) into (9), we come to a general answer for the heat flux and temperature profile in a system with an arbitrary mass distribution. While these integrals cannot be performed in a general case, they permit a number of conclusions.

To begin with, it turns out that a flow in a system depends only on the boundary masses and the maximum mass in the chain (which specifies the minimum frequency). We readily see that if the masses are constant, we come to the standard answer for the heat flow [3]. In the reverse approximation, when $M_1 \gg M_0$, we obtain:

$$J = \frac{2\sqrt{2}\Delta T}{\pi\omega_0\gamma^2}\left(\frac{\phi(0)}{\phi(1)}\right)^2\left(\sqrt{2}\omega_{min}\gamma - \arctan(\sqrt{2}\omega_{min}\gamma)\right)$$

Up to this point, we assumed the boundary masses not to depend on N. We can include this dependence in the case of linearly increasing masses. Setting $M_1 \sim M_0 N^s$, the heat flux will acquire the form:

$$J = \frac{8\Delta T}{3\pi}\omega_0^2\gamma\frac{1}{N^{3s/2}}.$$

It is easy to verify the correctness of the assumptions we have made here, because we know the exact solution for the linear case.

As follows from the expression for the flux J, for a constant gradient $s = 1$ the system behaves as a thermal insulator, $J \sim N^{-3/2}$, and for $s = 0$ the flux does not depend on system size. In the intermediate case of $s = 2/3$, the system conforms to Fourier's law $J \sim N^{-1}$. Note that the dependence of heat flux on the size of a harmonic system is specified by boundary conditions [12], and it is possible to select them such that Fourier's law in the system will be obeyed. In the $s = 2/3$ case we selected appropriately the system itself while leaving unchanged standard boundary conditions. Thus a system with linearly increasing masses can be adjusted such that it will conform to Fourier's law.

Let us turn now to the temperature profile. First, as seen from relations (14), if the masses are not constant within the chain, the temperature at the points at which $M'(a) = 0$ should rise strongly.

Second, it can be shown that in the case of constant masses thermal profile within the system $T(a) = T$ is constant, a point well enough known. In the reverse approximation, in the case of linearly increasing masses, when $M_1 \gg M_0$, we come to:

$$T_a = T - \frac{1}{\pi}\Delta T \arcsin\left(\sqrt{a}\right).$$

We see immediately that the temperature in the system is determined by the more massive end. Now at the center of the system under consideration a temperature gradient $-\Delta T/\pi$ will appear.

In conclusion, we have presented the solution to systems of linear equations (2) describing systems of functional-gradient materials. In particular, we analyzed the problem of a nonequilibrium steady state of a harmonic variable-mass system connected with heat reservoirs which are maintained at different temperatures.

We have shown that in the particular case of linearly increasing masses the heat flux depends on system size $J \sim 1/N^{3s/2}$, with the exponent in this relation being determined by the law governing the increase of the boundary mass $M \sim N^s$. This result finds ready explanation when we turn to Fig. 2. The systems under consideration maintain vibrations of two types, delocalized phonons which transport heat ballistically and localized "gradons" . By properly varying the boundary mass as a function of N, we modulate in this way the number of phonons, to arrive finally at the dependence of the heat flux J on the system size N. A similar effect accounts for the appearance of a thermal gradient in a system with linearly increasing masses. We have demonstrated in the particular example of a system with linearly increasing masses that a harmonic system can both sustain formation of a temperature gradient and conform to Fourier's law.

I wish to express gratitude to A. Ya. Vul' for the assistance he has been rendering throughout my work on the paper, as well as to K. Yu. Platonov for our discussions which assisted in formulation of this problem.

---

[*] Ioffe Physical-Technical Institute, 194021 St Petersburg, Polytechnicheskaya 26, Russia; Reich@mail.ioffe.ru

[1] A. Dhar, Advances in Physics **57**, 457 (2008).

[2] V. Kannan, A. Dhar, and J. L. Lebowitz, Phys. Rev. E **85**, 041118 (2012).

[3] Z. Rieder, J. L. Lebowitz, and E. Lieb, Journal of Mathematical Physics **8**, 1073 (1967).

[4] J. J. Xiao, K. Yakubo, and K. W. Yu, Phys. Rev. B **73**, 054201 (2006).

[5] T. N. Shah and P. Gajjar, Physics Letters A **376**, 438 (2012).

[6] E. P. Jiao Wang and G. Casati, Phys. Rev. E **86**, 010101 (2012).

[7] J. Cheh and H. Zhao, Journal of Statistical Mechanics: Theory and Experiment **2011**, P10031 (2011).

[8] E. Pereira, Phys. Rev. E **83**, 031106 (2011).

[9] G. N. Watson, *A Treatise on the Theory of Bessel Functions Cambridge Mathematical Library* (Cambridge University Press; 2 edition, 1995) p. 814.

[10] Y. Yan and H. Zhao, Journal of Physics: Condensed Matter **24**, 275401 (2012).

[11] P. E. Hopkins and J. R. Serrano, Phys. Rev. B **80**, 201408 (2009).

[12] A. Dhar, Phys. Rev. Lett. **86**, 5882 (2001).

[13] K. Yakubo, J. J. Xiao, and K. W. Yu, Journal of Physics: Conference Series **92**, 012008 (2007).